



A Survey of Giardia and Cryptosporidium.....

A SURVEY OF GIARDIA AND CRYPTOSPORIDIUM SPP. IN RURAL
AND URBAN COMMUNITY IN NORTH DELTA, EGYPT

Khaled Abd El-Aziz Mohammad¹ & Alaa Abd El-Aziz Mohammad²

Parasitology Department, Faculty of medicine (Damietta)¹

Zoology Department, Faculty of science(Cairo)² Al-Azhar University)

ABSTRACT

Cryptosporidium spp. and *Giardia duodenalis* are 2 protozoan parasites that affect humans and a wide range of domestic and wild animals. These parasites are a major cause of diarrheal disease in humans and animals worldwide, causing high morbidity in their hosts, in immunocompromised hosts and children, they can lead to death. In order to assess the prevalence of these protozoans among population in the northern part of Delta and the risk of human infection, 330 people (175 females and 155 males) with or without symptoms who attended a Health Care Center were parasitologically studied. The purpose of this study was to assess the frequency of *Cryptosporidium* spp. and *Giardia* spp. in the northern part of Delta (Damietta governorate) and the risk of human infection of these protozoa related to personal data, socio-cultural and environmental characteristics, and the presence of symptoms/signs among a rural and urban population. Of the surveyed population (330), 3.6% were infected with *Cryptosporidium* spp. only or 24.2% with *Giardia* spp. and 2.4% were infected with both protozoa. The frequency of infection according to sex; 2.7% of males were infected and 0.9% of females by *Cryptosporidium* spp, 12.7% and 11.5% by *Giardia* spp. and 1.5% and 0.9% by both parasites, respectively. *Cryptosporidium* spp. and *Giardia* spp. was frequent in children (under 12 year) than adults in both sexes, however, *Cryptosporidium* infections were most frequently diagnosed in children less than 12years old and prevalence decreased with age. The prevalences of these protozoa in this community are lower than those reported by other studies, which is probably associated with the low density of the studied population. Statistical analysis revealed that a male sex, children, loss of weight, impropal waste disposal, diarrhea and abdominal pain were correlated with the presence of these parasites, which indicate the importance of these factors in rural communities.

Keywords: *Cryptosporidium* spp., *Giardia* spp, epidemiology, rural and urban community, North Delta, Damietta Governorate.

Corresponding author : Alaa abdel-Aziz Mohammed e-mail: alaasamn@yahoo.com mobile:0106780563

INTRODUCTION

Cryptosporidium spp. and *Giardia duodenalis* are 2 protozoan parasites that affect humans and a wide range of domestic and wild animals [1-3]. These parasites are a major cause of diarrheal disease in humans and animals worldwide, causing high morbidity in their hosts, and in immunocompromised hosts, they can lead to death. The transmission of these 2 parasites is sustained by zoonotic (animal to human) and anthropontic (human to human) cycles

where several species and genotypes are enrolled [3-5]. The protozoan flagellate *Giardia lamblia* shares several epidemiologic characteristics with *C. parvum*. Both organisms are water-borne and this route has been the cause of like *C. parvum*, *G.lamblia* is a cause of childhood diarrhea and may be transmitted by close-contact and is often associated with epidemics of diarrhea in day care facilities. 16 The prevalence of both *G. lamblia* and *C. parvum* is generally higher among very young children and this may be

A Survey of Giardia and Cryptosporidium.....

related to more efficient fecal-oral transmission of the infective stages or enhanced susceptibility due to lack of immunity.¹⁷ While the prevalence of cryptosporidiosis and giardiasis is well documented in patients and children with human immunodeficiency virus (HIV) and AIDS, data are not widely available on the cross-sectional distribution of these infections. Water-borne outbreaks of infections with the parasites have highlighted the fact that morbidity may occur outside of these focus groups in endemic areas. The infected hosts shed in the environment a large number of these transmissible and infective stages, contributing to an increase of environmental contamination. Cysts and oocysts not only remain infective for long periods in environment but are also resistant to the conventional treatment processes of water, representing a serious problem of public health [2,8-12]. This problem is also potentiated by the fact that the number of parasites required to induce infection is small, i.e., infectious dose of 83-123 oocysts for *Cryptosporidium* spp. and 19-50 cysts for *G. duodenalis* [1,13]. intestinal infections caused by *Cryptosporidium* and *Giardia* spp. are associated to climatic factors, basic sanitary conditions, and sociocultural characteristics [7]. Traditionally, the prevalence of these infections was considered a "stable endemic characteristic" resulting from the dynamics of repeated re-infection [1]. (According to Atías[1] and[7], the frequency at which repeated infection occurs in a population depends on the "infection pressure" and host susceptibility. "Infection pressure" is defined as the number of host exposures at a given location over a certain period of time. The factors involved in "infection pressure" are of two types: environmental and socio-economic, which are intimately linked since the environment has a certain influence on the socio-

economic conditions of a population and human activities also affect the environment [15]. Environmental conditions (e.g. temperature, humidity, wind, soil) and socio-economic factors are responsible for spreading and for developing infectious forms of *Cryptosporidium* and *Giardia* spp., and for polluting the environment with these agents. Giardiasis is a cosmopolitan zoonotic infection, and predominates in mild humid climates. According to Crompton [4], two hundred million are infected with *Giardia* spp. worldwide. And, 15% of the rural population in Latin America is estimated to be infected with these parasites; with a higher prevalence among children [1]. In addition, the infection prevalence is lower in the upper strata of the population, and increases in day care centers attended by the children of poorer families [6]. On the other hand, *Cryptosporidium* is an intestinal parasite that infects both in humans and animals. Infection with *Cryptosporidium* spp. was reported in immunocompromised, as well as immunocompetent, children and adults [1] and in animals [15]. The disease occurs in most countries including Arabian countries [19], USA [9] and Egypt [1, 8, 9, 15&19]. It also has a higher prevalence in tropical regions, reaching infection levels of 20-50%, especially in rural areas [9]. Moreover, both protozoan infections are associated with non-specific gastrointestinal symptoms [10] diarrhea and weight loss. [3]. The purpose of this study was to assess the frequency of *Giardia* spp. and *Cryptosporidium* in the northern part of Delta and the risk of human infection of these protozoa related to personal data, socio-cultural and environmental characteristics, and the presence of symptoms/signs among a rural and urban population.

Subjects, Materials and Methods
Study area

A Survey of Giardia and Cryptosporidium.....

North Delta, Damietta Governorate, Damietta city representing urban area and Kafr saad villages representing rural area (10km apart from Damietta). Damietta is a governorate situated in the north region of Delta, which lies in northeast Egypt some

200 km from the capital city, Cairo ([Fig. 1](#)). Million five hundred thousand people reside in an area of 589 km² (991,687 inhab rural areas). The town has a commercial downtown or "urban region", surrounded by a rural area.

Fig. 1: Geographical location of North Delta of Egypt.

**Selected population**

The study extended from January 2010 to January 2011, stool samples were referred to Department of parasitology, Faculty of medicine(Damietta), Al-Azhar University, Egypt Total 330 people (175 females and 155 males), with or without symptoms, who attended a Health Care Center were studied. Of these, 191 people lived in an urban area (UA) (57.4%) and 139 in a rural area (RA) (42.6%); 239 were under 12 years old, and 134 lived in the UA. Each individual was surveyed for personal, environmental and socio-cultural data. Personal information - age, sex, and place of residence - was collected, as was socio-cultural and environmental information, garbage disposal (open pit, burning, burying or city garbage collection), water supply (running water, home pump, community pump), tap (inside/outside, public), body waste disposal (sewage, well with a chamber, cesspool, or latrine), overcrowding (presence or absence, presence was defined as more than three people sleeping to a room), and house flooding (never, sometimes, frequent). In addition, subjects were asked about the presence or absence of the following symptoms/signs: anal itching, abdominal

pain, sleeping disorders, diarrhea, vomiting, appetite loss, weakness, and weight lost

Parasite analysis

A serial parasitological analysis was performed, consisting of stool sampling from a daily spontaneous bowel movement, sample were stored in a jar containing 5% formaldehyde. To detect *Giardia* spp. and *Cryptosporidium*, the samples were processed and subjected to direct wet smear method and Sheather's sugar flotation and acid-ether sedimentation techniques Technique [22]. Processed pellets were stained with lugol and modified Ziehl-Neelsen acid fast stain and observed under a microscope .

Statistical analysis


Significant associations were determined using the chi square test and Fisher's test. Socio-cultural and environmental characteristics, symptoms, age, and sex were used as (independent) explanatory variables. SPSS software, version 11.5 was used throughout.

Results


Fig (1) show detecting Cyst of *G. lamblia* in iodine preparation (A) and *Cryptosporidium*

A Survey of Giardia and Cryptosporidium.....

oocysts in Midified ZN acid fast stain



preparation (B)



Sex, age, and place of residence in the analyzed population are shown in Table 1. In the studied population, 3 % (10/330) of subjects were infected with *Cryptosporidium* spp. only, 22.7% (75/330) with *Giardia* spp. only and 4.5% (15/330) were infected with both protozoa. Regarding *Cryptosporidium* spp. infection, 5.6% (9/158) of males were

infected and 1.7% (3/172) of females, similarly for *Giardia* spp., 26.5% (42/158) of males were infected and 22% (38/172) of females, and 3.1% (5/158) and 2.9% (5/172) were infected with both protozoa. These apparent differences between the sexes were not significant.

Tab 1: age, Sex and place of residence of the 330 people studied in North Delta, Damietta

Areas	Sex	Under 12 year		Over 12 year	
		No (%)	No (%)	No (%)	No (%)
Urbun	Male	78(23.6)		19(5.7)	
	Female	61(18.5)		33(10)	
	Total	139(42.1)		52(15.7)	
Rural	Male	45(13.6.1)		16(4.8)	
	Female	55(16.6)		23(6.9)	
	Total	100(30.3)		39(11.8)	

The prevalence of *Cryptosporidium* spp. was 3.1% (6/191) in the UA and 4.0% (8/139) in the RA. *Giardia* spp. prevalence was similar in both areas, i.e., 23.5% (45/191) in the UA and 21.5% (30/139) in the RA. Of the 10 people parasitized by both protozoa, 5 (2.6%) lived in the UA and 5 (3.5%) in the RA. Distributions according age, sex, and place of residence are shown in Table 2. Analysis of the age distributions of both parasites in the populations revealed that the

frequency of *Cryptosporidium* spp. infection was highest among those aged 2 to 7 years, peaking at 4 years of age. On the other hand, *Giardia* spp. less age dependent and was found to be present in subjects up to 70 years old, peaking between the ages 4 and 12. When the data of both parasite infections were evaluated, mean age for *Cryptosporidium* spp. infection was 7 years and for *Giardia* spp. infection 24 to 35 years.

A Survey of Giardia and Cryptosporidium.....
Table 2: Frequencies of *Cryptosporidium* spp. And *Giardia* spp. infections in the Damietta population, with respect to sex, place of residence and age

Parasites	Males				Females			
	Under 12 year		Over 12 year		Under 12 year		Over 12 year	
	RA %	UA %	RA %	UA %	RA %	UA %	RA %	UA %
<i>Cryptosporidium</i> spp	3(6.6)	6(7.6)	0(0.0)	0(0.0)	3(5.4)	0(0.0)	0(0.0)	0(0.0)
<i>Giardia</i> spp	11(24.4)	22(28.2)	5(31.2)	4(21)	10(18.1)	16(26.2)	4(17.3)	8(24.2)
<i>Cryptosporidium+Giardia</i>	0(0.0)	3(3.8)	1(6.2)	1(5.2)	2(3.6)	0(0.0)	1(4.3)	0(0.0)

Association between clinical manifestation of *Cryptosporidium* and *Giardia lamblia* infection with socio-cultural and environmental data of the study population When these variables and the presence/absence of symptoms/signs were correlated with parasite frequency are shown in Table 3.

Tab 3: Socio-cultural and environmental variables of the 330 study subjects

Occupation	Education	Socioeconomic		Housing& Waste disposal		Water supply		Clinical symptoms			
		No. (%)	No. (%)	No. (%)	No. (%)	No. (%)	No. (%)	No. (%)	No. (%)		
House wife	95 (28.7)	Illiterate	135 (40.9)	Low	145 (43.9)	Modern building	135 (40.9)	Piped	60 (18.1)	Asympto -matic	198 (60)
Farmer	75 (22.7)	Primary school	95 (22.7)	Moderate	130 (39.3)	Primitiv e building	195 (59)	River nile	95 (28.7)	Diarrhea	36 (10.9)
Manual worker	75 (22.7)	preparatory school	65 (19.6)	High	55 (16.6)	Overcr- od	235 (71.2)	Shallo- w wells	105 (31.8)	Weight loss	15 (4.5)
Students	55 (16.6)	Secondary school	35 (10.6)		Sewage disposal	155 (46.9)	Deep wells	70 (21.2)	Abd. Pain	57 (17.2)	
Educated (High)	30 (9.0)	Above second			Well with chamber disposal	175 (53)			Nausea & Vomiting	24 (7.2)	

DISCUSSION

Intestinal parasites are very common in developing countries and *Cryptosporidium* and *Giardia* have been revealed as one of the most common parasites. *Cryptosporidium* sp. is a primary pathogen causing acute diarrhea and the most evident symptom of cryptosporidiosis is diarrhea. Non-specific

signs such as dehydration, fever, anorexia, weakness, and progressive loss of weight may be accompanied. Diarrhea is usually self-limiting in immunocompetent humans; however, it can be life-threatening in children and immunocompromised humans. Its prevalence varies according to different regions [5] . In developed countries, massive

A Survey of Giardia and Cryptosporidium.....

Cryptosporidium food-borne and water-borne outbreaks have been reported, while in developing countries, *Cryptosporidium* is associated with significant morbidity and mortality, especially among infants and children[27] . In Egypt, human cryptosporidiosis been reported with variable prevalence rates by [1, 8, 9, 15&19] The overall frequency of *Giardia* spp. infection found in the present study was higher than those reported for similar studies in Italy (3.5%,[13] ; but it is lower than those reported in Venezuela (28%,[16] , Chennai (26%, [11]), Pakistan (28.78%, in [23], and in Brazil (27.4%,[18] . Similar results were obtained in studies carried out in Thailand (5.3%,[28] , and Spain (5.05%,[20] .Other studies carried out in Argentina have reported higher rates of these parasitizes. Thus, Gamboa et al. [12] reported 34% for a shantytown, Guignard et al. [14] 23% in a population living in temporary homes, and Borda et al. [4] who reported 29% in shantytown. As regards studies carried out on the prevalence of *Cryptosporidium* in similar populations in other countries, our results are higher than those published by Wang et al. [29] in China (1% and 3.96%), Windsor et al. [31] in the United Kingdom (2.9%),[6] in Spain (1.5%),[28] in Thailand (0.8%), and [17] in Venezuela (1.6%). Percentages similar to those of the present study were found by [26] in members of the Thai Army and their families (4.1%). As regards data collected in Argentina, [12] report 48% in a shantytown, [14] 44.4% among those living in temporary homes, [21] 65.8% among a suburban population, and [24] 24.6%. A compared with previous Argentinean figure our prevalence rates for both parasites are notably lower. This may be due to the rural nature of our study population, and socio-economic, cultural, and demographic (fewer inhabitants per square km) differences, which would be expected reduce infection pressure.

Regarding sex, these protozoa were markedly more prevalent among males in the UA group, with an outstanding higher frequency in under 12-year olds. This is probably because the direct transmission of these parasites is related to higher population densities in the urban area, and because of the different hygiene habits of males and females in the same setting, which is reflected by the relatively low/high prevalence observed in the infant population. The category less than 12 years old was a risk factor for this infection. These results agree with the development of acquired immunity to both parasites consequent to repeated infections in childhood [10]. Of the socio-cultural and environmental variables studied, frequent flooding, dirt floor inside the home, cardboard/tin homes, and body waste disposal by latrine were associated with the prevalence of these parasites. and/or blastocystis. Therefore, the presence of these conditions could be used to estimate the risk of infection by these parasites. These data reinforce the existing strong relation between deficient sanitary conditions and intestinal parasitism, as has been mentioned by several Argentinean authors[4,12&24]. Several authors reported a correlation between diarrhea, abdominal pain, flatulence, itching, vomiting, or appetite loss, and the presence of both protozoan infections[6&17]. In this study only abdominal pain was found to be significantly associated with the presence of *Cryptosporidium* and/or *Giardia* spp. The present study is one of the first to associate the prevalence of *Giardia* spp. and/or *Cryptosporidium* with socio-cultural and environmental conditions, and with the clinical features of a North Delta population. We conclude that the low population density of the area studied substantially creates a lower infection pressure, and explains the lower infection prevalence's of the present study. The risk factors for acquiring *G.* spp

A Survey of Giardia and Cryptosporidium.....

and *C. spp* in Damietta are apparently similar and this is reflected in their age-prevalence profiles. Alternately, age-related acquisition of immunity to infection or loss of infection may be different for the parasites and this may contribute to the observed patterns that most infections with the parasites were in children, and clinical diagnosis was not an accurate indicator of infection. Factors such as malnutrition and bottle feeding as seen in some Rural areas may be important in determining the epidemiology of the infections in North Delta. The prevalence of *G. spp.* infections increased with age and this may be reflective of the increasing exposure as children attend schools and day-care facilities. Perhaps feeding or infection from untreated water may be more important for transmission of *C. spp* in young children.

REFERENCES

1. Abaza, S.M.; Maklouf, L.; El Shwey, Kh. And El Moamiiy, A.A.(1995): Intestinal opportunistic parasites among the different groups of immunocompromised hosts. J. Egypt. Soc. Parasitol., 25(3): 713-727.
2. Atías A. El hospedero. La relación hospedero-parasito. In: Atias A, editor. Parasitología Médica. Mediterraneo, Santiago, Chile: 1999. pp. 49–53.
3. Atwill, E.R. and Kemp, Z.S.(1998): Prevalence and associated risk factors for shedding cryptosporidium oocysts and cyst within fecal populations in calefornia App. Environment. Microbial. 63: 3446-49.
4. Borda CE, Rea MJ, Rosa JR, Maidana C. Intestinal parasitism in San Cayetano, Corrientes, Argentina. Bull Pan Am Health Organ. 1996;30:227–233. [PubMed]
5. Caccio SM, Ryan U. Molecular epidemiology of giardiasis. Mol Biochem Parasitol. 2008;160:75–80
6. Carbajal JA, Villar L, Lanuza MD, Esteban JG, Munoz C, Borras R. Clinical significance of *Blastocystis hominis* infection: epidemiologic study. Med Clin (Barc) 1997;108:608–612.
7. Crompton DW. The public health importance of hookworm disease. Parasitology. 2000;121:S39–S50.
8. El Mansoury ST, Abou El Naga IF, Negm AY, Amer EE. Influence of temperature and salinity on the viability and infectivity of *Giardia lamblia* and *Cryptosporidium parvum*. J Egypt Soc Parasitol. 2004;34:161–172. [PubMed]
9. El Shazly, A.M.; Soltan, D.M.; El-Sheikha, H.M.; Sadek, G.S. and Morsy, A.T.A. (2007): Correlation of ELISA coproantigen and oocysts count to the severity of cryptosporidiosis parvum in children. J. Egypt. Soc. Parasitol., 37(1): 107-120.
10. Faubert G. Immune response to *Giardia duodenalis*. Clin Microbiol Rev.2000;13:35–54.
11. Fernández MC, Veghese S, Bhuvaneswari R, Elizabeth SJ, Matheu T, Anitha A, Chitra AK. A comparative study of the intestinal parasites prevalent among children living in rural and urban settings in and around Chennai. J Commun Dis. 2002;34:35–39.
12. Gamboa MI, Basualdo JA, Cordoba MA, Pezzani BP, Minvielle MC, Lahitte HB. Distribution of intestinal parasitoses in relation to environmental and sociocultural parameters in La Plata, Argentina. J Helminthol. 2003;77:15–20.
13. Giacometti A, Cirioni O, Fortuna M, Drenaggi D, Vecchia S, Dérrico MM, Calisse G. Giardiasis: a parasitic disease of continued topicality. Study of prevalence among a selected adult population. Infez Med. 2000;8:82–86.
14. Guignard S, Aruebtu HM, Freyre L, Lujan H, Rubinstein H. Prevalence of enteroparasites in a residence for children in the Córdoba Province, Argentina. Eur J Epidemiol. 2000;16:287–293.
15. Mahmoud, Kh .A. (2006): Zoonotic cryptosporidiosis in man and animal in farms, Giza Governorate, Egypt. J Egypt Soc Parasitol, 36(2, Suppl.): 49-58
16. Miller SA, Rosario CL, Rojas E, Scorza JV. Intestinal parasitic infection and associated symptoms in children attending day care centres in Trujillo, Venezuela. Trop Med Int Health. 2003;8:342–347.

A Survey of Giardia and Cryptosporidium.....

17. Miller WA, Lewis DJ, Pereira MD, Lennox M, Conrad PA, Tate KW, Atwill ER. Farm factors associated with reducing Cryptosporidium loading in storm runoff from dairies. *J Environ Qual.* 2008;37:1875–1882.
18. Newman RD, Moore SR, Lima AA, Nataro JP, Guerrant RL, Sears CL. A longitudinal study of Giardia lamblia infection in north-east Brazilian children. *Trop Med Int Health.* 2001;6:624–634.
19. Noureldin,M.S.A.; Shaltout, A.A.; El Hunshary, E.M. and Ali, M.E. (1999): Opportunistic intestinal protozoa infections in immunocompromised children. *J. Egypt. Soc. Parasitol.*, 29(3): 951-961.
20. Perez Armengol C, Ariza Astolfi C, Ubeda Ontiveros JM, Guevara Benitez DC, de Rojas Alvarez M, Lozano Serrano C. Epidemiology of children's intestinal parasitism in the Guadalquivir Valley, Spain. *Rev Esp Salud Publica.* 1997;71:547–552.
21. Pezzani BC, Minvielle MC, De Luca MM, Radman N, Iacoy P, Basualdo Farjat JA. Survey of intestinal parasites in a periurban community from the province of Buenos Aires, Argentina. *Bol Chil Parasitol.* 1996;51:42–45.
22. Sheather, A. L. 1923. The detection of intestinal protozoan and mange parasites by a flotation technique. *J. Comp. Pathol.* 36:266–275.
23. Siddiqui MI, Bilquees FM, Hiyas M, Perveen S. Prevalence of parasitic infections in a rural area of Karachi, Pakistan. *J Pak Med Assoc.* 2002;52:315–320.
24. Soriano SV, Barbieri LM, Pierangeli NB, Giayetto AL, Manacorda AM, Castronovo E, Pezzani BC, Minvielle MC, Basualdo JA. Intestinal parasites and the environment: Frequency of intestinal parasites in children of Neuquen, Patagonia, Argentina. *Rev Latinoam Microbiol.* 2001;43:96–101.
25. Stenzel DJ, Boreham PF. *Blastocystis hominis* revisited. *Clin Microbiol Rev.* 1996;9:563–584.
26. Taamasri P, Mungthin M, Rangsin R, Tongupprakarn B, Areekul W, Leelayoova S. Transmission of intestinal blastocystosis related to the quality of drinkingwater. *Southeast Asian J Trop Med Public Health.* 2000;31:112–117.
27. Xiao L, Fayer R. Molecular characterisation of species and genotypes of *Cryptosporidium* and *Giardia* and assessment of zoonotic transmission. *Int J Parasitol.* 2008;38:1239–1255. [[PubMed](#)]
28. Waikagul J, Krudsood S, Radomyos P, et al. A crosssectional study of intestinal parasitic infections among schoolchildren in Nan Province, Northern Thailand. *Southeast Asian J Trop Med Public Health.* 2002;33:218–223.
29. Wang KX, Li CP, Wang J, Cui YB. Epidemiologic survey of *Blastocystis hominis* in Huainan City, Anuhui Province, China. *World J Gastroenterol.* 2002;8:928–932.
30. Wilson, M.E. (1991): *A World Guide to Infections: Diseases, Distribution, Diagnosis*, New York, Oxford, Oxford University Press
31. Windsor JJ, Macfarlane L, Hughes-Thapa G, Jones SK, Whiteside TM. Incidence of *Blastocystis hominis* in faecal samples submitted for routine microbiological analysis. *Br J Biomed Sci.* 2002;59:154–157