

KETAMINE PLUS NITROGLYCERIN VERSUS KETAMINE AS ADJUNCTS TO LIDOCAINE FOR INTRAVENOUS REGIONAL ANESTHESIA IN ADULT PATIENTS UNDERGOING HAND SURGERY

Islam A. Eliwa MD

Department of Anesthesiology, Faculty of Medicine, Zagazig University

Corresponding author:

Islam A. Eliwa

Tel. 01288778805

ABSTRACT

Background: Different agents have been used as adjuncts to lidocaine for intravenous regional anesthesia (IVRA) to improve the quality of analgesia.

Objective: The aim of this double blinded randomized controlled trial is to evaluate if adding nitroglycerin to ketamine is more effective than using ketamine alone as an adjuvant to lidocaine for IVRA.

Methods: Fifty adult ASA I-II patients undergoing hand surgery using IVRA were divided into two equal groups: ketamine/nitroglycerin (K/N) group received lidocaine 3mg/kg (maximum 200 mg) with ketamine 0.1 mg/kg (maximum 10 mg) plus nitroglycerin 2 mcg/kg (maximum 200 mcg) and ketamine (K) group received lidocaine 3 mg/kg (maximum 200 mg) with ketamine 0.1 mg/kg (maximum 10 mg). Statistical analysis was done using Student's t test or Chi square analysis whenever appropriate.

Results: The two groups were similar in demographic data and patients characteristics. The onset times of sensory and motor blockades were significantly shorter in K/N group. Although the preoperative visual analogue scale for pain intensity (VAS) was comparable, the intra-operative and postoperative VAS, were significantly less in K/N group. The intra-operative fentanyl consumption as an intra-operative analgesia was significantly less in K/N group. Also, the time to call for the first postoperative analgesia was significantly longer in K/N group than K group. However the postoperative analgesics consumption showed no significant difference between the two groups.

Conclusion: In comparison to ketamine, adding a combination of ketamine and nitroglycerin to lidocaine for IVRA improved the quality of anesthesia without inducing significant clinical side effects.

Key words: Nitroglycerin, ketamine, intravenous, regional, anesthesia.

INTRODUCTION

Ketamine hydrochloride acts as a non-competitive antagonist for N-methyl-D-aspartate (NMDA) receptors.^[1] Therefore, ketamine found a place in the field of IVRA as an effective additive,^[2] or even in higher doses as a sole agent with local anesthetic qualities.^[3] Nitroglycerin as a nitric oxide generator has been proved in several studies to have analgesic enhancement effects in acute and chronic pain conditions.^{[4],[5]} Recently, it was investigated as an adjuvant to lidocaine for IVRA with promising results.^[6]

The present double blinded randomized controlled trial aimed to compare between a combination of ketamine plus nitroglycerin versus ketamine solely as adjuncts to lidocaine for IVRA.

METHODS

After approval of local ethical committee and getting an informed written consent we

carried out this study at Zagazig university hospitals from January 2014 to December 2016. Fifty, ASA I-II, adult patients, age from 21 to 60 years of both genders who were undergoing hand surgery which was expected to be less than one hour duration, were included in the study. Exclusion criteria were decompensated cardiovascular or respiratory disease, sickle cell anemia, Reynaud's disease, peripheral cyanosis or neuropathy, regular use of analgesics, drug abuse and known allergy to one of the examined drugs.

All patients were pre-medicated with 2 mg midazolam and atropine 0.01 mg/kg half an hour before operation. Non invasive monitoring of electrocardiography, blood pressure and oxygen saturation were established. Two cannulae were inserted and secured: one in a vein on the dorsum of the operated hand for injection of the local anesthetic and the investigated

additives and the second cannula in the other hand for intravenous fluids and other drugs administration. A double cuffed pneumatic tourniquet was applied to the limb to be operated upon after exsanguinations with an Esmarch bandage. The proximal cuff was inflated to 100 mmHg above patient's systolic pressure and circulatory isolation of the arm was verified by absence of radial pulse and loss of the pulse oximeter trace in the ipsilateral index finger. Then, patients were randomly assigned into one of two groups with twenty five patients in each group: Ketamine/Nitroglycerin (K/N) group in which patients received lidocaine 3 mg/kg (maximum 200 mg) plus ketamine 0.1 mg/kg (maximum 10 mg) and nitroglycerine 2 mcg/kg (maximum 200 mcg) versus Ketamine (K) group whereas patients received lidocaine 3 mg/kg (maximum 200 mg) and ketamine 0.1 mg/kg (maximum 10 mg). The drugs used in either group were pre-prepared, mixed and diluted together up to 40 ml with normal saline by an anesthesiologist who was not included in the anesthesia administration. On the other hand, the patient and the anesthesiologist who provided and monitored the local intravenous anesthesia intra and postoperatively didn't know to which group the patient was allocated. When complete sensory and motor block was achieved, the distal cuff was inflated to 250 mmHg and the proximal cuff was deflated and surgeon was asked to proceed in the procedure to be done. At the end of operation, the tourniquet was released by cyclic deflation technique. It was not permitted the release of tourniquet before thirty minutes or keeping it inflated more than one hour and half.

Outcome measurements:

- Onset time of sensory block which was defined as the time elapsed from injection of the study drugs to the loss of sensation to pinprick in dermatomes of radial, median and ulnar nerves.

- Onset time of motor block which was defined as the time elapsed from injection of the study drugs to the inability of patient to flex the wrist or fingers.

- Visual analogue scale (VAS) for evaluation of pain intensity (zero value is the best and ten value is the worst) was measured one hour preoperatively, at thirty minutes after tourniquet inflation and fifteen minutes postoperatively after complete deflation of the tourniquet and before administration of any postoperative analgesia.

- Heart rate and blood pressure were recorded at five minutes intervals and any increases or decreases by 25% of the baseline were reported.

- Intra-operative analgesic requirements were estimated, hence 1 mcg/kg of fentanyl was given on patient request or with VAS > than 4 value.

- Time to the first call for analgesia which was defined as the time elapsed from the complete deflation of the tourniquet at the end of operation until the patients asked for postoperative analgesia.

- Postoperative consumption of analgesics whereas patients received ketorolac 30 mg IM on patients request for analgesia with a minimum 6 hours interval and a maximum dose 120 mg/24 hours.

- Side effects such as delirium, loss of consciousness and local anesthetic toxicity were noted.

Statistical analysis: The collected data from patients were verified prior to computerized data entry. The Statistical Package for Social Sciences (SPSS for Windows, Version 19) was used for statistical analysis. Simple randomization was done using random number generator in SPSS. All data are presented as mean \pm standard deviation or number (percentage). Statistical analysis was done using Student's t test or Chi square analysis whenever appropriate. All tests of significance were two-tailed with *P* values < 0.05 were considered statistically significant. Based on a priori sample size calculations that incorporated results from

previous studies,^[7] it was estimated that a twenty five patients per group (total fifty patients) would be required for a statistical

power of 80 % at a two tailed level of significance of $P < 0.05$.

RESULTS

Table (1): Demographic data and patients characteristics.

	Ketamine/Nitroglycerin (K/N) group (N = 25)	Ketamine (K) group (N = 25)	
Age (year)	41 ± 8.4	38 ± 10	0.26
Weight (kg)	81 ± 6.7	83 ± 7.3	0.36
Height (cm)	178 ± 3.8	175 ± 5.9	0.09
Sex:			0.38
- Male	14 (56)	17 (68)	
- Female	11 (44)	8 (32)	
Type of surgery:			0.85
- Carpal tunnel	12 (48)	9 (42)	
- Tendon repair	6 (24)	7 (28)	
- Contracture release	4 (16)	5 (20)	
- Trigger finger	3 (12)	4 (16)	
Duration of surgery (min)	59 ± 10	57 ± 9	0.41

- Data are presented as mean ± standard deviation or number (percentage).

Table (2): The onset times of sensory and motor blockades.

	Ketamine/Nitroglycerin (K/N) group (N = 25)	Ketamine (K) group (N = 25)	
Onset time of sensory block (min)	3.7 ± 0.57 *	4.2 ± 0.63	0.004
Onset time of motor block (min)	5.3 ± 0.64 *	6.1 ± 0.87	0.001

- Data are presented as mean ± standard deviation.

- * Significant $P < 0.05$.

Table (3): preoperative, intra-operative and postoperative visual analogue scale (VAS) for pain intensity in the operated arm.

	Ketamine/Nitroglycerin (K/N) group (N = 25)	Ketamine (K) group (N = 25)	
Preoperative VAS	0.8 ± 0.81	0.6 ± 0.74	0.59
Intra-operative VAS	1.8 ± 1.1 *	2.7 ± 1.3	0.01
Postoperative VAS	2.1 ± 1.06 *	2.8 ± 1.05	0.02

- Data are presented as mean ± standard deviation.

- VAS: Visual analogue scale for pain intensity with (0) value is the best and (10) value is the worst pain encountered.

- * Significant $P < 0.05$.

Table (4): Intra-operative and postoperative analgesics consumption.

	Ketamine/Nitroglycerin (K/N) group (N = 25)	Ketamine (K) group (N = 25)	
Intra-operative analgesia:			
- Patients required rescue fentanyl (number of patients)	3 (12) *	9 (36)	0.04
- Consumed amount of fentanyl (mcg)	10 ± 27 *	30 ± 41	0.04
Time to call for first postoperative analgesia (min)	67 ± 17 *	50 ± 18	0.002
Postoperative analgesia:			
- consumed ketorolac (mg)	64 ± 20	74 ± 24	0.14

- Data are presented as mean ± standard deviation or number (percentage).

- * Significant P < 0.05.

The two groups were similar in demographic data and patients characteristics including age, weight, height, gender, type of operation and duration of surgery (table 1). The onset times of sensory and motor blockades were significantly shorter in the K/N group (table 2). Although, the preoperative VAS was comparable, the intra-operative and postoperative VAS, were significantly less in K/N group than K group (table 3). As been shown in (table 4), the intra-operative fentanyl consumption as an intra-operative rescue analgesic was significantly less in the K/N group. Also, the time to call for the first postoperative analgesia was significantly longer in the K/N group than the K group. However, the postoperative analgesics consumption showed no significant difference between the two groups. No patient in the study developed hemodynamic increases or decreases by 25% of baseline blood pressure and heart rate or other clinical side effects were noted.

DISCUSSION

The problematic issue with IVRA is the application of a tourniquet that should be kept inflated all the time of surgical

procedure to maintain the regional anesthesia and avoid local anesthetic toxicity.^[8] Also, one of its drawbacks is absence of postoperative analgesia after release of tourniquet.^[9] Several trials were carried out to find an additive to lidocaine,^{[10],[11],[12]} which promotes the relief of tourniquet pain, prolongs the duration of analgesia after tourniquet release and reduces the systemic analgesic drugs consumption.^[13]

Ketamine acts as a non-competitive NMDA receptors antagonist. NMDA receptors antagonists have been widely used for control of peri-operative pain.^[14] Hence the NMDA receptors have been found on peripheral unmyelinated sensory axons in addition to the spinal cord NMDA receptors, it can be explained why ketamine as an antagonist of NMDA receptors is capable to reduce tourniquet pain and improve the quality of anesthesia when added to lidocaine as an adjuvant in IVRA.^[15] However, patients exhibited disorientation and hallucinations after the release of tourniquet which limited the usefulness of ketamine for this purpose. These unpleasant effects were associated with ketamine use in doses ≥ 0.5 mg/kg of

body weight. When it was used in doses < 0.5 mg/kg in IVRA no central nervous system effects have been noticed.^[16] Therefore, in our study we used ketamine in a dose 0.1 mg/kg to avoid these unpleasant side effects.

It is well known that nitroglycerin has a direct potent vasodilator effect that seems to enhance the distribution of lidocaine to nerves.^[17] This action of nitroglycerin is greatly dose dependent and can be increased by the increase in drug dose.^[18] Also, nitroglycerin exhibits its analgesic action because it is metabolized to nitric oxide in the cells.^[19] Nitric oxide elevates the intracellular concentration of cyclic guanosine monophosphate which modulates pain perception in the central and peripheral nervous systems.^[20] Additionally, nitroglycerin as a nitric oxide generator by topical application produces an anti-inflammatory effect and analgesia through blocking hyperalgesia and the neurogenic component of inflammatory oedema.^[21]

Asadi and Mehri,^[22] evaluated the analgesic effect of nitroglycerin when added to lidocaine on quality of intravenous regional anesthesia in patients undergoing elective forearm and hand surgery. Their study revealed that nitroglycerin as adjuvant drug to lidocaine is effective in improving the overall quality of anesthesia, shortening onset time of both sensory and motor blockades. Also, Elmetwaly et al,^[23] compared and evaluated the effect of adding ketamine or nitroglycerin to lidocaine for IVRA on intra-operative and postoperative analgesia, onset of sensory and motor blockade and tourniquet pain. Their conclusion was that either ketamine or nitroglycerin when added to lidocaine improved the overall quality of anesthesia, reduced the tourniquet pain and improved the postoperative analgesia consumption. Moreover, ketamine as an adjuvant produced better tolerance to tourniquet while nitroglycerin as an adjuvant

produced faster onset of sensory and motor blockades.

Our study evaluated if adding nitroglycerin to ketamine is more effective than using ketamine alone as an adjuvant to lidocaine for intravenous regional anesthesia. We found that the onset times of sensory and motor blockades were significantly shorter with adding nitroglycerin to ketamine. The tolerance of tourniquet pain was significantly better as reflected by lower intra-operative and postoperative VAS and less consumption of fentanyl as an intra-operative rescue analgesic. In addition, the time to call for the first postoperative analgesia was significantly longer with using the combination of nitroglycerin and ketamine. However, the postoperative analgesic consumption was similar in both groups.

Despite both ketamine and nitroglycerin are expected to cause tachycardia, our results did not show any significant hemodynamic changes or clinical side effects. This can be attributed to the small doses of both drugs which were used in the study (ketamine 0.1 mg/kg with maximum dose 10 mg, nitroglycerin 2 mcg/kg with maximum dose 200 mcg). Also, the presence of tourniquet, avoidance of tourniquet deflation before 30 min and the use of cyclic deflation at the end of operation allowed gradual escape of both drugs to systemic circulation which limited their hemodynamic effects.^[24] However, one of the limitations of the present study is the small number of patients which was enough to detect significant clinical differences between the two groups but is difficult to detect significant side effects. Because side effects are not common it necessitates a study of a huge number of patients to assure their statistical significance.

CONCLUSION

Adding a combination of ketamine and nitroglycerin to lidocaine for intravenous regional anesthesia improved the quality of anesthesia, shortened the onset times of sensory and motor blockade and reduced

the intra-operative analgesic consumption without inducing significant clinical side effects.

REFERENCES

- De Kock MF, Lavand'homme PM. The clinical role of NMDA receptor antagonists for the treatment of postoperative pain. *Best Pract Res Clin Anaesthesiol* 2007; 21: 85 – 98.
- Durrani Z, Winnie AP, Zsigmond EK, Burnett ML. Ketamine for intravenous regional anesthesia. *Anesth Analg* 1989; 68: 328 – 332.
- Kaul TK, Mittal BR, Gupta S. comparison of lignocaine and ketamine as intravenous regional anaesthetics. *Indian J Anaesth* 1993; 41: 234 – 239.
- Sen S, Ugur B, Aydin ON, Ogurlu M, Gursoy F, Savk O. The analgesic effect of nitroglycerin added to lidocaine on intravenous regional anesthesia. *Anesth Analg* 2006; 102 (3): 916 – 920.
- Lauretti GR, Lima IC, Reis MP, Prado WA, Pereira NL. Oral ketamine and transdermal nitroglycerin as analgesic adjuvants to oral morphine therapy for cancer pain management. *Anesthesiology* 1999; 90:1528 – 1533.
- Honarmand A, Safavi M, Fatemy A. The analgesic effect of three different doses of nitroglycerine when added to lidocaine for intravenous regional anesthesia in trauma patients. *Turkish journal of Trauma and Emergency surgery* 2011; 17: 497 – 503.
- Kumar A, Sharma DK, Datta B. Addition of ketamine or dexmedetomidine to lignocaine in intravenous regional anesthesia: A randomized controlled study. *J anaesthesiol Clin Pharmacol* 2012; 28 (4): 501 – 504.
- Mohr B. safety and effectiveness of intravenous regional anesthesia (Bier block) for outpatient management of forearm trauma. *CJEM* 2006; 8 (4): 247 – 250.
- Jankovic RJ, Visnjic MM, Milic DJ, Stojanovic MP, Djordjevic DR, Pavlovic MS. Does the addition of ketorolac and dexamethasone to lidocaine intravenous regional anesthesia improve postoperative analgesia and tourniquet tolerance for ambulatory hand surgery? *Minerva Anesthesiol* 2008; 74: 521 – 527.
- Memis D, Turan A, Karamanlioglu B, Pamukcu Z, Kurt Z. Adding Dexmedetomidine to lidocaine for intravenous regional anesthesia. *Anesth Analg* 2004; 98: 835 – 840.
- Rayan AA, El Sayed AA. Efficacy of intravenous regional anesthesia with dexmedetomidine: local addition versus systemic infusion. *Ain-Shams J Anaesthesiol* 2016; 9: 83 – 91.
- Nagpal V, Rana S, Singh J, Chaudhary SK. Comparative study of systemically and perineurally administered tramadol as an adjunct for supraclavicular brachial plexus block. *Journal of Anaesthesiology Clinical pharmacology* 2015; 31(2): 191 – 195.
- Choyce A, Peng P. A systematic review of adjuncts for intravenous regional anesthesia for surgical procedures. *Can J Anaesth* 2002; 49 (1): 32 – 45.
- Kurdi MS, theerth DA, Deva RS. Ketamine: Current applications in anesthesia, pain and critical care. *Anesthesia Essays and Researches* 2014; 8 (3): 283 – 290.
- Viscomi CM, Friend A, Parker C, Murphy T, Yarnell M. Ketamine as an adjuvant in lidocaine intravenous regional anesthesia: a randomized, double-blind, systemic control trial. *Reg Anesth Pain Med* 2009; 34: 130 – 133.
- Gorgias NK, Maidatsi PG, Kyriakidis AM, Karakoulas KA, Alvanos DN, Giala MM. Clonidine versus ketamine to prevent tourniquet pain during intravenous regional anesthesia with lignocaine. *Reg Anesth Pain Med* 2001; 26: 512 – 517.
- Abbasivash R, Hassani E, Aghdashi M, Shirvani M. The effect of nitroglycerin as an adjuvant to lidocaine in intravenous regional anesthesia. *Anesth* 2009; 20 (2): 265 – 269.
- Hashimoto S, Kobayashi A. Clinical pharmacokinetics and pharmacodynamics of glyceryl trinitrate and its metabolites. *Clin Pharmacokinet* 2003; 42: 205 – 221.
- Abu Elyazed M, Abdel-Ghaffar M. Nitroglycerin and ketorolac as an adjuvant to lidocaine for intravenous regional anesthesia in patients undergoing forearm and hand surgery. *AAMJ* 2014; 12 (4):159 – 176.
- Kumar R, Biyani G, Mohmmmed S, Karnawat R, Sethi P, Sharma V. Efficacy of nitroglycerin as an adjuvant to intravenous regional anesthesia: a review of literature.

- Indian journal of clinical anaesthesia 2015; 2 (2): 107 – 110.
21. Lauretti GR, Perez MV, Reis MP, Pereira NL. Double-blind evaluation of transdermal nitroglycerin as adjuvant to oral morphine for cancer pain management. *J Clin Anesth* 2002; 14: 83 – 86
22. Asadi HK, Mehri D. The analgesic effect of nitroglycerin added to lidocaine on quality of intravenous regional anesthesia in patients undergoing elective forearm and hand surgery. *Acta Cir. Bras.* 2013; 28 (1): 19-25.
23. Elmetwaly KF, Hegazy NA, Aboelseoud AA, Alshaer AA. Does the use of ketamine or nitroglycerin as an adjuvant to lignocaine improve the quality of intravenous regional anesthesia? *Saudi J Anaesth* 2010; 4: 55 – 62.
24. Barazandeh N, Ebtahaj M, Naderi F. The effect of adding nitroglycerin as an adjuvant to lidocaine on pain and hemodynamic variables of intravenous regional anesthesia upper extremity surgery. *J Mazand Univ Med Sci* 2014; 24 (109): 55 – 61.